arXiv:math/0606716v1 [math.AG] 28 Jun 2006

REDUCTION METHOD FOR LINEAR SYSTEMS OF PLANE CURVES WITH BASE FAT POINTS

MARCIN DUMNICKI

Abstract. In the paper we develop a new method of proving non-speciality of a linear system with base fat points in general position. Using this method we show that the Hirschowitz-Harbourne Conjecture holds for systems with base points of equal multiplicity bounded by 42.

1. Introduction

Let $\mathbb{K}$ be a field of characteristics zero, $\mathbb{N} = \{0, 1, 2, \dots\}$, $\mathbb{N}^* = \{1, 2, \dots\}$, let $\mathbb{R}$ denote the field of real numbers.

Definition 1. Let $D \subset \mathbb{N}^2$ be finite (any such set will be called a *diagram*), let $m_1, \dots, m_r \in \mathbb{N}^*$, let $p_1, \dots, p_r \in \mathbb{K}^2$. Define the $\mathbb{K}$-vector space $\mathcal{L}_D(m_1p_1, \dots, m_rp_r) \subset \mathbb{K}[X, Y]$:

$$\mathcal{L}_D(m_1p_1, \dots, m_rp_r) := \Big\{ f = \sum_{(\beta_1,\beta_2)\in D} c_{(\beta,\beta_2)} X^{\beta_1} Y^{\beta_2} \mid c_{(\beta_1,\beta_2)} \in \mathbb{K}, \frac{\partial^{\alpha_1+\alpha_2} f}{\partial X^{\alpha_1} \partial Y^{\alpha_2}}(p_j) = 0, \alpha_1 + \alpha_2 < m_j, j = 1, \dots, r \Big\}.$$

Define the *dimension* of the *system of curves* $\mathcal{L}_D(m_1, \dots, m_r)$ to be

$$\dim \mathcal{L}_D(m_1, \dots, m_r) := \min_{\{p_j\}_{j=1}^r, p_j \in \mathbb{K}^2} \dim_{\mathbb{K}} \mathcal{L}_D(m_1p_1, \dots, m_rp_r) - 1.$$

Remark 2. If points $p_1, \dots, p_r$ are in general position we have

$$\dim \mathcal{L}_D(m_1, \dots, m_r) = \dim_{\mathbb{K}} \mathcal{L}_D(m_1p_1, \dots, m_rp_r) - 1.$$

The system $\mathcal{L}_D(m_1, \dots, m_r)$ can be understood as a vector space of curves generated by monomials with exponents from D having multiplicities at least $m_1, \dots, m_r$ in r general points.

Definition 3. Let $L = \mathcal{L}_D(m_1, \dots, m_r)$ be a system of curves. Define the *virtual dimension* vdim *of* L and the *expected dimension* edim *of* L

$$\operatorname{vdim} L := \#D - 1 - \sum_{j=1}^{r} \binom{m_j + 1}{2},$$

$$\operatorname{edim} L := \max\{\operatorname{vdim} L, -1\}.$$

1991 *Mathematics Subject Classification.* 14H50; 13P10.

Key words and phrases. Multivariate interpolation, linear systems, Hirschowitz–Harbourne conjecture.

Intuitively, we should have $\dim L = \operatorname{edim} L$.

Definition 4. We say that a *system of curves L is special* if

$$\dim L > \operatorname{edim} L.$$

Otherwise we say that L *is non-special*.

Observe that by linear algebra we have always $\dim L \geq \operatorname{edim} L$ since multiplicity m imposes $\binom{m+1}{2}$ conditions.

2. The Hirschowitz–Harbourne Conjecture

For systems of the form $\mathcal{L}_d(m_1, \ldots, m_r) := \mathcal{L}_D(m_1, \ldots, m_r)$, $D = \{\alpha \in \mathbb{N}^2 \mid |\alpha| \leq d\}$ the well-known Hirschowitz–Harbourne Conjecture giving geometrical description of the speciality of a system was formulated in [9]. To formulate this conjecture consider the blowing-up $\pi : \widetilde{\mathbb{P}}^2 \to \mathbb{P}^2$ in r general points with exceptional divisors $E_1, \ldots, E_r$.

Definition 5. A curve $C \subset \mathbb{P}^2$ is said to be a -1-*curve* if it is irreducible and the self-intersection $\widetilde{C}^2$ of its proper transform $\widetilde{C} \subset \widetilde{\mathbb{P}}^2$ is equal to -1.

Conjecture 6 (Hirschowitz–Harbourne)**.** *A system $L = \mathcal{L}_d(m_1, \ldots, m_r)$ is special if and only if there exists a -1-curve $C \subset \mathbb{P}^2$ such that*

$$\widetilde{L}.\widetilde{C} \leq -2,$$

where $\widetilde{L} := |d\pi^*(\mathcal{O}_{\mathbb{P}^2}(1)) - \sum_{j=1}^{r} m_j E_j|$.

This conjecture was studied by many authors, we refer only to the recent results. For homogenous systems ($m_1 = m_2 = m_3 = \cdots = m_r =: m$), the above conjecture holds for $m \leq 20$ (see [4, 5]). In the general case the Conjecture holds for multiplicities bounded by 11 (see [8]). Both these results were obtained with the help of computers.

For further information about the above conjecture see for example [3].

In this paper we develop a new method (called "diagram cutting") based on some properties of matrices assigned to systems of curves. This method allows providing easy proofs of non-speciality for large family of systems. Moreover, these proofs can often be found argorithmically with a computer program. Sometimes this needs a lot of computations, but then it can easily be checked "by hand".

As a result of the method we show that in order to check non-speciality of all homogeneous systems of bounded multiplicity m it is enough to check a finite number of cases. This result was obtained in purely theoretical way.

The second result is Theorem 31 stating that Conjecture 6 holds for homogeneous multiplicities bounded by 42. This result was obtained using a computer program.

3. Diagram cutting method

Before introducing the method we must establish the notation and say when a system is non-special in the language of matrices.

Definition 7. Let $j \in \mathbb{N}^*$, $\alpha \in \mathbb{N}^2$. Define the mapping

$$\varphi_{j,\alpha} : \mathbb{K}[X,Y] \ni f \mapsto \frac{\partial^{|\alpha|} f}{\partial X^\alpha}(P_{j,X}, P_{j,Y}) \in \mathbb{K}[P_{j,X}, P_{j,Y}],$$

where $P_{j,X}, P_{j,Y}$ are new indeterminates used instead of X, Y.

We will use the following notation: $\mathcal{M}(n,k;R)$ will denote the set of all matrices with n rows, k columns, and coefficients from R (a ring or a field). For an $M \in \mathcal{M}(n,k;R)$ we write $M_{[j,\ell]}$ for the element of M in the j-th row and the ℓ-th column.

Definition 8. Let $L = \mathcal{L}_D(m_1, \ldots, m_r)$ be a system of curves, let $D = \{(\alpha_{1,X}, \alpha_{1,Y}), \ldots, (\alpha_{n,X}, \alpha_{n,Y})\}$, $\alpha_{i,X}, \alpha_{i,Y} \in \mathbb{N}$ for $i = 1, \ldots, n$. Let $\mathcal{A} = \{(j,\beta) \in \mathbb{N} \times \mathbb{N}^2 \mid |\beta| < m_j,\ j = 1, \ldots, r\} = \{\mathfrak{a}_1, \ldots, \mathfrak{a}_c\}$. Define the matrix $M(L) \in \mathcal{M}(c, n; \mathbb{K}[P_{i,X}, P_{i,Y}]_{i=1}^r)$ as

$$M(L)_{[j,k]} := \varphi_{\mathfrak{a}_j}(X^{\alpha_{k,X}} Y^{\alpha_{k,Y}}).$$

For given points $p_1 = (p_{1,X}, p_{1,Y}), \ldots, p_r = (p_{r,X}, p_{r,Y}) \in \mathbb{K}^2$ we will use the natural evaluation mapping

$$\nu_{p_1,\ldots,p_r} : \mathbb{K}[P_{i,X}, P_{i,Y}]_{i=1}^r \ni f \mapsto f|_{P_{i,X} \mapsto p_{i,X}, P_{i,Y} \mapsto p_{i,Y}} \in \mathbb{K}.$$

Proposition 9. *Let $L = \mathcal{L}_D(m_1, \ldots, m_r)$ be a system of curves. Then* $\dim L = \#D - \operatorname{rank} M(L) - 1$.

Proof. Let $p_1, \ldots, p_r \in \mathbb{K}^2$ be points in general position. Consider the linear mapping

$$\Phi : \Big\{ f = \sum_{(\alpha_X, \alpha_Y) \in D} c_{(\alpha_X, \alpha_Y)} X^{\alpha_X} Y^{\alpha_Y} \Big\} \ni f \mapsto (\nu_{p_1,\ldots,p_r} \circ \varphi_{\mathfrak{a}_j}(f))_{j=1}^c \in \mathbb{K}^c.$$

We have $\mathcal{L}_D(m_1 p_1, \ldots, m_r p_r) = \ker \Phi$. Let M denote the matrix of Φ in bases $\{(\alpha_{1,X}, \alpha_{1,Y}), \ldots, (\alpha_{n,X}, \alpha_{n,Y})\}$, $\{\mathfrak{a}_1, \ldots, \mathfrak{a}_c\}$. We have

$$\nu_{p_1,\ldots,p_r}(M(L)_{[j,k]}) = M_{[j,k]},$$

hence $\operatorname{rank} M(L) = \operatorname{rank} M$ (we use the facts that $\operatorname{char} \mathbb{K} = 0$ and $p_1, \ldots, p_r$ are in general position). Now we compute

$$\dim L = n - \operatorname{rank} M - 1 = \#D - \operatorname{rank} M(L) - 1.$$

□

Definition 10. Define the bidegree $\operatorname{bdeg} : \mathbb{K}[P_{i,X}, P_{i,Y}]_{i=1}^r \to \mathbb{N}^2$ given by $\operatorname{bdeg}(P_{i,X}) := (1,0), \operatorname{bdeg}(P_{i,Y}) := (0,1)$ for $i = 1, \ldots, r$.

Proposition 11. *Let $\mathcal{L}_D(m_1,\ldots,m_r)$ be a system of curves. Let M be a square submatrix of $M(L)$ of size $s \in \mathbb{N}^*$. After a necessary renumbering of columns and rows we can assume that M is given by columns $(\alpha_{1,X},\alpha_{1,Y}),\ldots,(\alpha_{s,X},\alpha_{s,Y})$ and rows $\mathfrak{a}_1,\ldots,\mathfrak{a}_s$. Then $\det M$ is a bihomogeneous (w.r.t.* bdeg*) polynomial of bidegree $(\sum_{i=1}^s \alpha_{i,X}, \sum_{i=1}^s \alpha_{i,Y}) - \gamma$, where $\gamma \in \mathbb{N}^2$ depends only on the choice of rows.*

Proof.

$$\det M = \sum_{\sigma\in S_s} \operatorname{sgn}(\sigma) M_{[1,\sigma(1)]} \ldots M_{[s,\sigma(s)]}.$$

For $M_{[j,k]} \neq 0$ we have $\operatorname{bdeg} M_{[j,k]} = (\alpha_{k,X},\alpha_{k,Y}) - \beta_j$, where $\mathfrak{a}_j = (\ell_j,\beta_j)$ for some $\ell_j \in \mathbb{N}$, $\beta_j \in \mathbb{N}^2$. Hence

$$\operatorname{bdeg} M_{[1,\sigma(1)]} \cdots M_{[s,\sigma(s)]} = \left(\sum_{i=1}^s \alpha_{i,X}, \sum_{i=1}^s \alpha_{i,Y}\right) - \sum_{i=1}^s \beta_i.$$

We finish the proof by taking $\gamma = \sum_{i=1}^s \beta_i$. □

Proposition 12. *Let $m \in \mathbb{N}^*$, let $D \subset \mathbb{N}^2$. Assume that $\#D = \binom{m+1}{2}$. Then $L = \mathcal{L}_D(m)$ is non-special if and only if D do not lie on a curve of degree $m-1$.*

Proof. From the previous proof we can see that $\det M(L) = c \cdot f$, where f is a monomial, $c \in \mathbb{K}$. Let $D = \{(\alpha_{1,X},\alpha_{1,Y}),\ldots,(\alpha_{n,X},\alpha_{n,Y})\}$. For $\beta = (\beta_X,\beta_Y)$, $|\beta| < m$ we have

$$\begin{aligned} M(L)_{[(1,\beta),j]} &= \varphi_{(1,\beta)}(X^{\alpha_{j,X}} Y^{\alpha_{j,Y}}) \\ &= \prod_{k=1}^{\beta_X} (\alpha_{j,X} - k + 1) \cdot \prod_{k=1}^{\beta_Y} (\alpha_{j,Y} - k + 1) \cdot P_{1,X}^{\alpha_{j,X}-\beta_X} P_{1,Y}^{\alpha_{j,Y}-\beta_Y}. \end{aligned}$$

Since we are only interested in the value of c we would like to compute the determinant of $M = M(L)_{X\mapsto 1, Y\mapsto 1}$. By a suitable sequence of row operations we can change M into the following matrix M'

$$M'_{[(1,\beta),j]} = \alpha_{j,X}^{\beta_X} \alpha_{j,Y}^{\beta_Y}, \qquad \det M \neq 0 \iff \det M' \neq 0.$$

Now $\det M' = 0$ if and only if rows of M' are linearly dependent, but this happens if and only if D lies on a curve of degree $m-1$. □

Now we can present the diagram cutting method and prove that it can be used to bound the dimension of a system of curves.

Theorem 13. *Let $m_1,\ldots,m_r,m_{r+1},\ldots,m_p \in \mathbb{N}^*$, let $D \subset \mathbb{N}^2$ be a diagram, let $F : \mathbb{N}^2 \ni (\alpha_1,\alpha_2) \mapsto r_1\alpha_1 + r_2\alpha_2 + r_0 \in \mathbb{R}$ be an affine function, $r_0,r_1,r_2 \in \mathbb{R}$. Let*

$$\begin{aligned} D_1 &:= \{(\alpha_1,\alpha_2) \in D \mid F(\alpha_1,\alpha_2) < 0\}, \\ D_2 &:= \{(\alpha_1,\alpha_2) \in D \mid F(\alpha_1,\alpha_2) > 0\}. \end{aligned}$$

If $L_2 := \mathcal{L}_{D_2}(m_{r+1}, \ldots, m_p)$ *is non-special and* $\operatorname{vdim} L_2 = -1$ *then*

$$\dim \mathcal{L}_D(m_1, \ldots, m_p) \leq \dim \mathcal{L}_{D_1}(m_1, \ldots, m_r).$$

Proof. Put $L_1 := \mathcal{L}_{D_1}(m_1, \ldots, m_r)$. We can compute the dimension of a system $L := \mathcal{L}_D(m_1, \ldots, m_p)$ as $\dim L = \#D - \operatorname{rank} M(L) - 1$. As $D = D_1 \cup D_2$, in an appropriate basis the matrix $M(L)$ is of the following form

$$M(L) = \left[\begin{array}{c|c} M(L_1) & K_1 \\ \hline K_2 & M(L_2) \end{array}\right].$$

Take a maximal non-zero minor M' of $M(L_1)$ and consider the following square submatrix of $M(L)$:

$$M = \left[\begin{array}{c|c} M' & K_1' \\ \hline K_2' & M(L_2) \end{array}\right],$$

where K_1' and K_2' are suitable submatrices of K_1 and K_2. It suffices to show that $\det M \neq 0$. The columns of M' are indexed by elements from some $D_1' \subset D_1$ hence the columns of M are indexed by $D' := D_1' \cup D_2$. Let $U = [M'\ K_1']$, $L = [K_2'\ M(L_2)]$ be submatrices of M, let

$$\mathcal{C} := \{C \subset D' \mid \#C = \#D_2\}.$$

For $C \in \mathcal{C}$ define L_C (respectively U_C) as the submatrix of L (resp. U) given by taking the columns indexed by elements of C. Now we can compute

$$\det M = \sum_{C \in \mathcal{C}} \epsilon(C) \det L_C \det U_{D' \setminus C},$$

for $\epsilon(C) = \pm 1$. Observe that $\det L_C \in \mathbb{K}[P_{r+1,X}, P_{r+1,Y}, \ldots, P_{p,X}, P_{p,Y}]$, $\det U_{D' \setminus C} \in \mathbb{K}[P_{1,X}, P_{1,Y}, \ldots, P_{r,X}, P_{r,Y}]$ and consider $\det M$ as a polynomial of indeterminates $\{P_{\ell,X}, P_{\ell,Y}\}_{\ell > r}$ over $\mathbb{K}[\{P_{\ell,X}, P_{\ell,Y}\}_{\ell \leq r}]$. We have

$$\det M = \pm \det M' \det M(L_2) + f.$$

Assume that $\det M = 0$. As $\det M' \neq 0$ and $\det M(L_2) \neq 0$ from the assumptions, we must have another non-zero term $g \in \mathbb{K}[\{P_{\ell,X}, P_{\ell,Y}\}_{\ell > r}]$ appearing in f such that $\operatorname{bdeg}(g) = \operatorname{bdeg}(\det M(L_2))$. The term g was given by some $C \in \mathcal{C}$, $C \neq D_2$. Since

$$\operatorname{bdeg}(g) = \sum_{(\alpha_1, \alpha_2) \in C} (\alpha_1, \alpha_2) - \gamma, \quad \operatorname{bdeg}(\det M(L_2)) = \sum_{(\alpha_1, \alpha_2) \in D_2} (\alpha_1, \alpha_2) - \gamma$$

we have

$$F\Big(\sum_{(\alpha_1, \alpha_2) \in C} (\alpha_1, \alpha_2)\Big) = F\Big(\sum_{(\alpha_1, \alpha_2) \in D_2} (\alpha_1, \alpha_2)\Big).$$

As F is an affine form and $\#C = \#D_2$ we have

$$\sum_{(\alpha_1, \alpha_2) \in C} F(\alpha_1, \alpha_2) = \sum_{(\alpha_1, \alpha_2) \in D_2} F(\alpha_1, \alpha_2),$$

but from the definition of D_2 it is possible only when $C = D_2$, contradiction. □

4. Reduction of homogeneous systems

We will use the following notation for a sequence of multiplicities:

Definition 14. Let $m_1, \ldots, m_r \in \mathbb{N}^*$, $k_1, \ldots, k_r \in \mathbb{N}$. Define

$$(m_1^{\times k_1}, \ldots, m_r^{\times k_r}) = (\underbrace{m_1, \ldots, m_1}_{k_1}, \ldots, \underbrace{m_r, \ldots, m_r}_{k_r}).$$

We will use diagrams of the following form:

Definition 15. Let $a_1, \ldots, a_n, u_1, \ldots, u_n \in \mathbb{N}$. Define

$$(a_1^{\uparrow u_1}, \ldots, a_n^{\uparrow u_n}) := \bigcup_{i=1}^{n} \{i-1\} \times \{u_i, \ldots, u_i + a_i - 1\} \subset \mathbb{N}^2.$$

Example 16.




Fig. 1. Diagram $(2^{\uparrow 3}, 1^{\uparrow 0})$




Fig. 2. Diagram $(2^{\uparrow 3}, 1^{\uparrow 0}, 0^{\uparrow 0}, 3^{\uparrow 2})$

Observe that $\#(a_1^{\uparrow u_1}, \ldots, a_n^{\uparrow u_n}) = \sum_{i=1}^{n} a_i$.

Definition 17. Let $\mathbb{Z}$ denote the set of all integers. We say that two diagrams D_1, D_2 are *equivalent* (or D_1 *is equivalent to* D_2) if there exists $\alpha \in \mathbb{Z}^2$ such that $D_1 = D_2 + \alpha$.

Remark 18. Observe that diagram $(0^{\uparrow 0}, \ldots, 0^{\uparrow 0}, a_1^{\uparrow u_1}, \ldots, a_n^{\uparrow u_n})$ is equivalent to $(a_1^{\uparrow u_1}, \ldots, a_n^{\uparrow u_n})$ is equivalent to $(a_1^{\uparrow u_1 + u}, \ldots, a_n^{\uparrow u_n + u})$.

Proposition 19. *Let $m_1, \ldots, m_r \in \mathbb{N}^*$, let D_1, D_2 be diagrams. If D_1 and D_2 are equivalent, then* $\dim \mathcal{L}_{D_1}(m_1, \ldots, m_r) = \dim \mathcal{L}_{D_2}(m_1, \ldots, m_r)$.

Proof. Let $D_1 + (\alpha_1, \alpha_2) = D_2$, let $p_1, \ldots, p_r \in \mathbb{K}^2$ be points in general position. The maps

$$\mathcal{L}_{D_1}(m_1 p_1, \ldots, m_r p_r) \ni f \mapsto X^{\alpha_1} Y^{\alpha_2} f \in \mathcal{L}_{D_2}(m_1 p_1, \ldots, m_r p_r),$$
$$\mathcal{L}_{D_2}(m_1 p_1, \ldots, m_r p_r) \ni f \mapsto X^{-\alpha_1} Y^{-\alpha_2} f \in \mathcal{L}_{D_1}(m_1 p_1, \ldots, m_r p_r)$$

are well-defined (we can assume that none of the coordinates of $p_1, \ldots, p_r$ are zero), linear and inverse to each other. □

Lemma 20. *Let $m \in \mathbb{N}^*$, $D = (1^{\uparrow m-1}, 2^{\uparrow m-2}, \ldots, (m-1)^{\uparrow 1}, m^{\uparrow 0})$. Then $\mathcal{L}_D(m)$ is non-special and* $\operatorname{vdim} \mathcal{L}_D(m) = -1$.

Proof. By Lemma 12 it is enough to show that D (Fig. 3 shows an example for $m = 3$) do not lie on a curve C of degree $m-1$. Let $L_j = \{(x, y) \in \mathbb{R}^2 \mid x + y + j = 0\}$, $j = 0, \ldots, m-1$. Observe that $\#(D \cap L_j) = j + 1$ so by Bezout Theorem and induction we have $\bigcup_{j=0}^{m-1} L_j \subset C$, a contradiction. □

Remark 21. Observe that we can do the same for diagram $(m^{\uparrow 0}, (m-1)^{\uparrow 0}, \ldots, 1^{\uparrow 0})$.

Lemma 22. *Let $m \in \mathbb{N}^*$, $D = (m^{\uparrow m}, m^{\uparrow m-1}, \ldots, m^{\uparrow 0})$. Then $\mathcal{L}_D(m^{\times 2})$ is non-special and* $\operatorname{vdim} \mathcal{L}_D(m^{\times 2}) = -1$.

Proof. Let $F = y - m + \frac{1}{2}$. Observe that $D = D_1 \cup D_2$ (Fig. 4 shows an example for $m = 3$), where

$$D_1 := \{p \in D \mid F(p) < 0\} = (0, 1^{\uparrow m-1}, 2^{\uparrow m-2}, \ldots, m-1^{\uparrow 1}, m^{\uparrow 0}),$$
$$D_2 := \{p \in D \mid F(p) > 0\} = (m^{\uparrow m}, m-1^{\uparrow m}, \ldots, 1^{\uparrow m}).$$

The diagram D_1 is equivalent to $(1^{\uparrow m-1}, 2^{\uparrow m-2}, \ldots, (m-1)^{\uparrow 1}, m^{\uparrow 0})$ hence from Lemma 20 the system $\mathcal{L}_{D_1}(m)$ is non-special. The diagram D_2 is equivalent to $(m^{\uparrow 0}, (m-1)^{\uparrow 0}, \ldots, 1^{\uparrow 0})$ so $\mathcal{L}_{D_2}(m)$ is non-special. As $\#D_2 = \binom{m+1}{2}$, we can use Theorem 13 to obtain non-speciality of $\mathcal{L}_D(m^{\times 2})$. □

Lemma 23. *Let $m, k \in \mathbb{N}^*$, $D = (m^{\uparrow k-1}, m^{\uparrow k-2}, \ldots, m^{\uparrow 0})$. If $m+1|k$ then $L = \mathcal{L}_D(m^{\times 2\frac{k}{m+1}})$ is non-special and* $\operatorname{vdim} L = -1$.

Proof. We proceed by induction on k. For $k = m+1$ we use the previous Lemma. Let $k > m+1$. Take $F = x-(m+1)+\frac{1}{2}$. Observe that $D = D_1 \cup D_2$ (Fig. 5 shows an example for $m = 3$, $k = 12$), where

$$D_1 := \{p \in D \mid F(p) < 0\} = (m^{\uparrow k-1}, \ldots, m^{\uparrow k-1-m}),$$
$$D_2 := \{p \in D \mid F(p) > 0\} = (0, \ldots, 0, m^{\uparrow k-(m+1)-1}, \ldots, m^{\uparrow 0}).$$

The diagram D_1 is equivalent to $(m^{\uparrow m}, \ldots, m^{\uparrow 0})$, hence from Lemma 22 the system $\mathcal{L}_{D_1}(m^{\times 2})$ is non-special. The diagram D_2 is equivalent to $(m^{\uparrow k-(m+1)-1}, \ldots, m^{\uparrow 0})$ and from the induction hypothesis we get that system $\mathcal{L}_{D_2}(m^{\times 2\frac{k-(m+1)}{m+1}})$ is non-special. Now, Theorem 13 allows us to finish the proof. □

Lemma 24. *Let $m, k, h \in \mathbb{N}^*$, $D = (h^{\uparrow k-1}, h^{\uparrow k-2}, \ldots, h^{\uparrow 0})$. If $m+1|k$ and $m|h$ then $L = \mathcal{L}_D(m^{\times 2\frac{kh}{m(m+1)}})$ is non-special and* $\operatorname{vdim} L = -1$.

Proof. We proceed by induction on h. If $h = m$ then the situation was treated in the previous Lemma. Let $h > m$. Take $F = y+x-(k-1+m)+\frac{1}{2}$. Observe that $D = D_1 \cup D_2$ (Fig. 6 shows an example for $m = 3$, $k = 12$, $h = 9$), where

$$D_1 := \{p \in D \mid F(p) < 0\} = (m^{\uparrow k-1}, m^{\uparrow k-2}, \ldots, m^{\uparrow 0}),$$
$$D_2 := \{p \in D \mid F(p) > 0\} = ((h-m)^{\uparrow k-1+m}, \ldots, (h-m)^{\uparrow m}).$$

According to Lemma 23 the system $\mathcal{L}_{D_1}(m^{\times 2\frac{k}{m+1}})$ is non-special. The diagram D_2 is equivalent to $((h-m)^{\uparrow k-1}, (h-m)^{\uparrow k-2}, \ldots, (h-m)^{\uparrow 0})$ and by

the induction hypothesis the system $\mathcal{L}_{D_2}(m^{\times 2\frac{k(h-m)}{m(m+1}})$ is non-special. Again we finish the proof by using Theorem 13. □

Definition 25. Let $m \in \mathbb{N}^*$, $h = m(m+1)$. Define the set (called the *end of layer systems*)

$$\mathrm{EoLS}(m) = \{\mathcal{L}_D(m^{\times 2k+h-1}) \mid D = (h^{\uparrow k-1}, \ldots, h^{\uparrow 0}, (h-1)^{\uparrow 0}, \ldots, 1^{\uparrow 0}), k = 1, \ldots, m+1\}.$$

Observe that for every $L \in \mathrm{EoLS}(m)$ we have $\mathrm{vdim}\, L = -1$.

Lemma 26. *Let $m, k \in \mathbb{N}^*$, $h = m(m+1)$, $p = 2k+h-1$, $D = (h^{\uparrow k-1}, h^{\uparrow k-2}, \ldots, h^{\uparrow 0}, (h-1)^{\uparrow 0}, \ldots, 1^{\uparrow 0})$. If the set* $\mathrm{EoLS}(m)$ *contains only non-special systems then the system $L = \mathcal{L}_D(m^{\times p})$ is non-special.*

Proof. Take $k_1, k_2 \in \mathbb{N}$ such that $k = k_1(m+1) + k_2$, $1 \le k_2 \le m+1$. Put $F = x - k_1(m+1) + \frac{1}{2}$. Observe that $D = D_1 \cup D_2$ (Fig. 7 shows an example for $m = 3$, $k = 11$), where

$$D_1 := \{p \in D \mid F(p) < 0\} = (h^{\uparrow k-1}, h^{\uparrow k-2}, \ldots, h^{\uparrow k_2}),$$
$$D_2 := \{p \in D \mid F(p) > 0\} = (h^{\uparrow k_2-1}, \ldots, h^{\uparrow 0}, h-1^{\uparrow 0}, \ldots, 1^{\uparrow 0}).$$

The diagram D_1 is equivalent to the diagram $(h^{\uparrow k-k_2-1}, h^{\uparrow k-k_2-2}, \ldots, h^{\uparrow 0})$ and since $m+1 | k-k_2$, it follows from Lemma 24 that $\mathcal{L}_{D_1}(m^{\times 2\frac{(k-k_2)h}{m(m+1)}})$ is non-special. The system $\mathcal{L}_{D_2}(m^{\times 2k_2+h-1}) \in \mathrm{EoLS}(m)$ and we can use Theorem 13 to complete the proof. □

Theorem 27. *Let $m, d_L \in \mathbb{N}^*$. Assume that for $d = d_L, \ldots, d_L + m(m+1)$ every system $\mathcal{L}_d(m^{\times p})$, $p \in \mathbb{N}$, is non-special. Moreover, assume that the set* $\mathrm{EoLS}(m)$ *contains only non-special systems. Then for any $d \ge d_L, p \in \mathbb{N}$ the system $\mathcal{L}_d(m^{\times p})$ is non-special.*

Proof. We proceed by induction on d. For $d_L \le d \le d_L + m(m+1)$ the proof is obvious. Take $d > d_L + m(m+1)$. We want to show that the system $\mathcal{L}_D(m^{\times p})$ is non-special, where $D = ((d+1)^{\uparrow 0}, \ldots, 1^{\uparrow 0})$. Take $h = m(m+1)$, $F = y + x - (d-h) + \frac{1}{2}$. Observe that $D = D_1 \cup D_2$ (Fig. 8 shows an example for $m = 3$, $d_L = 3$, $d = 16$), where

$$D_1 := \{p \in D \mid F(p) < 0\} = (d+1-h^{\uparrow 0}, \ldots, 1^{\uparrow 0}),$$
$$D_2 := \{p \in D \mid F(p) > 0\} = (h^{\uparrow d+1-h}, \ldots, h^0, h-1^{\uparrow 0}, \ldots, 1^{\uparrow 0}).$$

As $d+1-h \ge d_L$ we may use the induction hypothesis for the system $\mathcal{L}_{D_1}(m^{\times p-(2d-h+3)})$. By Lemma 26 the system $\mathcal{L}_{D_2}(m^{\times 2d-h+3})$ is non-special. Again we finish the proof by using Theorem 13. □

Definition 28. Let $m, d_0 \in \mathbb{N}^*$. Put

$$S(m, d_0) := \mathrm{EoLS}(m) \cup$$
$$\{\mathcal{L}_d(m^{\times r}) \mid \operatorname{vdim} \mathcal{L}_d(m^{\times r}) \geq -2m^2, d_0 \leq d \leq d_0 + m(m+1), r \in \mathbb{N}^*\}.$$

Theorem 29. *Let $m, d_0 \in \mathbb{N}$. If the set $S(m, d_0)$ contains only non-special systems then every system $\mathcal{L}_d(m^{\times r})$ for $d \geq d_0$, $r \in \mathbb{N}$ is non-special.*

Proof. By Theorem 27 it suffices to show that every system $\mathcal{L}_d(m^{\times r})$, $r \in \mathbb{N}$, $d = d_0, \ldots, d_0 + m(m+1)$, is non-special. Let $L = \mathcal{L}_d(m^{\times r})$. If $\operatorname{vdim} L \geq -2m^2$ then L is non-special by the assumptions. Let $\operatorname{vdim} L < -2m^2$. From now on we use the notations and theory of reductions introduced in [7, 8]. We want to apply a sequence of r weak m-reductions to the diagram $D = \{\alpha \in \mathbb{N}^2 \mid |\alpha| \leq d\}$ to end with the empty diagram,

$$D \xrightarrow{m\,\mathrm{w}} D_1 \xrightarrow{m\,\mathrm{w}} D_2 \xrightarrow{m\,\mathrm{w}} D_3 \xrightarrow{m\,\mathrm{w}} \cdots \xrightarrow{m\,\mathrm{w}} \varnothing.$$

Following the notations of [8] let us consider a diagram $D = (a_1, a_2, \ldots, a_n)$. Observe that the m-reduction is not possible only if $a_i = a_{i+1} < m$. As D is the result of a sequence of weak m-reductions this can only happen for $i \leq 2m$. While performing a m-weak reduction we use at most m additional points for each a_i, $i = 1, \ldots, 2m$, and for each i it is sufficient to do it only once. So we use at most $2m^2$ additional points to reduce D to the empty diagram, hence if $\operatorname{vdim} L < -2m^2$ then L is non-special. □

Example 30.




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8

5. Homogeneous systems with bounded multiplicity

Theorem 31. *The Hirschowitz-Harbourne Conjecture holds for homogeneous systems with multiplicities bounded by* 42.

Proof. For $m < 20$ the result can be found in [5]. For $m = 20, \ldots, 42$ we choose $d_0(m) = 3m$. In order to check the conjecture we have to do the following:

(1) Find all non-special systems among $\mathcal{L}_d(m^{\times r})$ for $d \leq d_0(m)$ (there are only finitely many of them). Next, for every such system we must show that it satisfies the Hirschowitz-Harbourne conjecture. This was done with the help of computer programs. By the proof of Theorem 29 the maximal size of matrix (for $m = 42$) can be 8128×11656, but in most cases the combination of reduction method and Cremona transformation gives an immediate answer.
(2) For every system from the set $S(m, d_0(m))$ we must prove its non-speciality.

As the set $S(m, d_0(m))$ contains systems with big size of diagrams, this cannot be done without preparations. For a system $L = \mathcal{L}_D(m^{\times r}) \in \mathrm{EoLS}(m)$ we use reduction method described in [8] to reduce the problem to the question about non-speciality of $L' = \mathcal{L}_{D'}(m^{\times 5})$ for some diagram D'. For $m = 42$ this forces us to compute the determinant of 4515×4515 matrix 43 times — this can be done. For the rest of the systems from $S(m, d_0(m))$ we use the following fact (see Proposition 28 in [8]).

Theorem 32. *Let $m_1, \ldots, m_r \in \mathbb{N}^*$. There exists a diagram D with the property: if $\mathcal{L}_D(m_1, \ldots, m_r)$ is non-special then for all $d \in \mathbb{N}$ the system $\mathcal{L}_d(m_1, \ldots, m_r)$ is non-special.*

So for each r such that $\mathcal{L}_d(m^{\times r}) \in S(m, d_0(m))$ we have to check only one system $\mathcal{L}_D(m^{\times r})$ for some diagram D depending on r. We also reduce this system to $\mathcal{L}_{D'}(m^{\times 9})$. In [6] the reader can find the table with the actual number of cases to be checked, as well as all necessary software with instructions, on how to perform the tests. □

Remark 33. The test decribed above can also be performed for greater values of m, but for each $m \geq 43$ this will take at least several days of computation. It seems that now it would be better to reorganize the method.

6. Closing remarks

Remark 34. There exists another method of proving non-speciality of given system (or for a family of systems) based on blowing-up the projective space introduced by C. Ciliberto and R. Miranda ([5]). It seems that diagram cutting method is different and sometimes works better. Moreover, all definitions and results of section 3 can be easily carried over to the higher-dimensional case of the systems of polynomials in n variables vanishing (with multiplicities) at points in general position. This is not known for the method of C. Ciliberto and R. Miranda.

Remark 35. Observe that Theorem 27 can be reformulated to the following.


Theorem 36. *If the set* EoLS(m) *contains only non-special systems and* $\mathcal{L}_d(m_1,\dots,m_r)$ *is non-special then*

$$\mathcal{L}_{d+m(m+1)}(m_1,\dots,m_r,m^{\times p})$$

is non-special, where $p=2d+m(m+1)+1$.

This shows that in order to find all non-special systems of the form $\mathcal{L}_d(m_1,\dots,m_r)$ with $m_i\le M$, $i=1,\dots,r$ it is sufficient to check a finite number of cases.

Example 37. We will show non-speciality of the system $L=\mathcal{L}_{21}(7^{\times 6},6^{\times 4},1)$ by diagram cutting method. The proof (found by a computer) can be easily read off from the picture. The system L was studied in [10].



References

[1] J. Alexander, A. Hirschowitz, *An asymptotic vanishing theorem for generic unions of multiple points*, Invent. Math. 140, 303–325 (2000)

[2] J. Alexander, A. Hirschowitz, *Polynomial interpolation in several variables*, J. Algebraic Geometry 4, 201–222 (1995).

[3] C. Ciliberto, *Geometric Aspects of Polynomial Interpolation in More Variables and of Waring's Problem*, Proc. ECM, Barcelona (2000).

[4] C. Ciliberto, F. Cioffi, R. Miranda, F. Orecchia, *Bivariate Hermite interpolation and linear systems of plane curves with base fat points*, Computer mathematics, Lecture Notes Ser. Comput., 10, World Sci. Publishing, River Edge, NJ, 87–102 (2003).
[5] C. Ciliberto, R. Miranda, *Linear systems of plane curves with base points of equal multiplicity*, Trans. Amer. Math. Soc. 352, 4037–4050 (2000).
[6] M. Dumnicki, `http://gamma.im.uj.edu.pl/dumnicki/interpol.htm`, June 1, 2006.
[7] M. Dumnicki, *Expected term bases for generic multivariate Hermite interpolation*, arXiv: math.AG/0503701.
[8] M. Dumnicki, W. Jarnicki, *New effective bounds on the dimension of a linear system in* $\mathbb{P}^2$, arXiv: math.AG/0505183.
[9] A. Hirshowitz, *Une conjecture pour la cohomologie des diviseurs sur les surfaces rationelles génériques*, J. Reine Angew. Math. 397, 208–213 (1989).
[10] S. Yang, *Linear systems in* $\mathbb{P}^2$ *with base points of bounded multiplicity*, arXiv:math.AG/0406591 v2.

Marcin Dumnicki
Institute of Mathematics, Jagiellonian University,
Reymonta 4, 30-059 Kraków, Poland